

YOU ARE CORDIALLY INVITED TO ATTEND  
A SPEAKER PROGRAM

# NERLYNX<sup>®</sup> (neratinib) Tablets for Early-Stage, HER2+ Breast Cancer

## PRESENTED BY:

BRIAN LEYLAND JONES, MD  
NATIONAL FOUNDATION FOR  
CANCER RESEARCH  
SIOUX FALLS, SD

## PROGRAM HOST:

JESSICA MAS  
415-350-3133

## PROGRAM INFORMATION:

SATURDAY, FEBRUARY 15, 2020  
8:05AM  
HILTON - HAWAIIAN VILLAGE  
TAPA BALLROOM  
2005 KALIA ROAD  
HONOLULU, HI 96815

## PLEASE REGISTER ONLINE AT:

[PUMAREG.TSGMEDED.COM](http://PUMAREG.TSGMEDED.COM)

Enter Event Code: 50723



New Jersey, Minnesota, Vermont, the Department of Defense, and the Department of Veterans Affairs have restrictions on receiving in-kind benefits (e.g., meals, paid parking) at company-sponsored events. For all attendees, please be advised that information such as your name and the value and purpose of any educational item, meal, or other items of value you receive may be publicly disclosed. You are accountable for understanding such restrictions and complying with them. If you are licensed in or affiliated with any of these states or federal agencies, Puma Biotechnology policies may restrict you from consuming any portion of the Puma Biotechnology-sponsored meal at this program or from receiving any other in-kind benefit from Puma Biotechnology (e.g., paid parking) in connection with the program.

This program is sponsored by Puma Biotechnology, Inc.

This is not an independent educational program, and no CME credits will be provided.

**Please see Important Safety Information on reverse side and accompanying  
Full Prescribing Information including Patient Information**

**nerlynx<sup>®</sup>**  
(neratinib) tablets

## IMPORTANT SAFETY INFORMATION

NERLYNX® (neratinib) tablets, for oral use

**INDICATIONS AND USAGE:** NERLYNX is a kinase inhibitor indicated for the extended adjuvant treatment of adult patients with early-stage HER2 overexpressed/amplified breast cancer, to follow adjuvant trastuzumab-based therapy.

**CONTRAINDICATIONS: None**

### WARNINGS AND PRECAUTIONS:

- **Diarrhea:** Aggressively manage diarrhea occurring despite recommended prophylaxis with additional antidiarrheals, fluids, and electrolytes as clinically indicated. Withhold NERLYNX in patients experiencing severe and/or persistent diarrhea. Permanently discontinue NERLYNX in patients experiencing Grade 4 diarrhea or Grade  $\geq 2$  diarrhea that occurs after maximal dose reduction.
- **Hepatotoxicity:** Monitor liver function tests monthly for the first 3 months of treatment, then every 3 months while on treatment and as clinically indicated. Withhold NERLYNX in patients experiencing Grade 3 liver abnormalities and permanently discontinue NERLYNX in patients experiencing Grade 4 liver abnormalities.
- **Embryo-Fetal Toxicity:** NERLYNX can cause fetal harm. Advise patients of potential risk to a fetus and to use effective contraception.

**ADVERSE REACTIONS:** The most common adverse reactions ( $> 5\%$ ) were diarrhea, nausea, abdominal pain, fatigue, vomiting, rash, stomatitis, decreased appetite, muscle spasms, dyspepsia, AST or ALT increase, nail disorder, dry skin, abdominal distention, weight decreased and urinary tract infection.

To report SUSPECTED ADVERSE REACTIONS, contact Puma Biotechnology, Inc. at 1-844-NERLYNX (1-844-637-5969) and [www.NERLYNX.com](http://www.NERLYNX.com) or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

### DRUG INTERACTIONS:

- Gastric acid reducing agents: Avoid concomitant use with proton pump inhibitors (PPI) and H<sub>2</sub>-receptor antagonists. Separate NERLYNX by 3 hours after antacid dosing. Separate NERLYNX by at least 2 hours before or 10 hours after H<sub>2</sub>-receptor antagonists.
- Strong or moderate CYP3A4 inhibitors: Avoid concomitant use.
- Strong or moderate CYP3A4 inducers: Avoid concomitant use.
- P-glycoprotein (P-gp) substrates: Monitor for adverse reactions of narrow therapeutic agents that are P-gp substrates when used concomitantly with NERLYNX.

### USE IN SPECIFIC POPULATIONS:

- Lactation: Advise women not to breastfeed.