

# Elevate the Possibilities: TRODELVY in Pretreated HR+/HER2- mBC and mTNBC

 **TRODELVY**<sup>®</sup>  
sacituzumab govitecan-hziy  
180 mg for injection

JOIN YOUR COLLEAGUES FOR AN EDUCATIONAL PROGRAM

## PROGRAM DESCRIPTION

Saturday, February 8, 2025

9:15 AM - 10:00 AM HT

The presenter will discuss key information for TRODELVY in HR+/HER2-negative mBC and mTNBC, including clinical trial data, safety, dosing, and more.

This is a non-CME/CE Session

## PRESENTED BY



**Chaitali Nangia, MD**  
Hematologist/Oncologist  
at Hoag Medical Group  
Co-Director, Hematology/Oncology  
at Chan Soon-Shiong Institute for Medicine  
Irvine, CA

## LOCATION

### 'Alohilani Resort Waikiki Beach

2490 Kalakaua Avenue  
Honolulu, Hawaii 96815  
Room: Bluefin I

**TRODELVY is a Trop-2-directed antibody and topoisomerase inhibitor conjugate indicated for the treatment of adult patients with:**

- Unresectable locally advanced or metastatic triple-negative breast cancer (mTNBC) who have received two or more prior systemic therapies, at least one of them for metastatic disease
- Unresectable locally advanced or metastatic hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative (IHC 0, IHC 1+ or IHC 2+/ISH-) breast cancer who have received endocrine-based therapy and at least two additional systemic therapies in the metastatic setting



**NCCN Category 1 | Preferred option for HR+/HER2- mBC**

**NCCN:** Sacituzumab govitecan-hziy (TRODELVY) is recommended as a Category 1 preferred treatment option for adult patients with recurrent unresectable or metastatic HR+/HER2- breast cancer who have received prior treatment, including endocrine therapy, a CDK4/6 inhibitor, and at least 2 lines of chemotherapy (including a taxane), at least one of which in the metastatic setting. It may be considered for later line if not used as second-line\* therapy.



**NCCN Category 1 | Preferred option for 2L and later mTNBC**

**NCCN:** Sacituzumab govitecan-hziy (TRODELVY) is recommended as a Category 1 preferred treatment option for adult patients with unresectable locally advanced or mTNBC who have received 2 or more prior systemic therapies, at least one of them for metastatic disease. It may be considered for later line if not used as second-line therapy.

NCCN makes no warranties of any kind whatsoever regarding their content, use, or application and disclaims any responsibility for their application or use in any way.

Category 1 indicates that based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.<sup>1</sup>  
\*If not a candidate for fam-trastuzumab deruxtecan-nxki.<sup>2</sup>

The NCCN recommendations may differ from the TRODELVY Prescribing Information.  
1L=first line; 2L=second line; CDK4/6=cyclin-dependent kinase 4/6; HER2=human epidermal growth factor receptor 2; HR+=hormone receptor-positive; IHC=immunohistochemistry; ISH=in situ hybridization; mBC=metastatic breast cancer; mTNBC=metastatic triple-negative breast cancer; NCCN=National Comprehensive Cancer Network.

## Gilead does not provide or reimburse for alcohol at Speaker Programs.

If you are a licensed US healthcare professional, the meal associated with this program may be reportable under the federal Open Payments/Sunshine Act and/or state law. Gilead policy and the PhRMA Code limit attendance at this promotional educational program to healthcare professionals. Attendance by guests cannot be accommodated.

## IMPORTANT SAFETY INFORMATION

### BOXED WARNING: NEUTROPENIA AND DIARRHEA

- **Severe or life-threatening neutropenia may occur. Withhold TRODELVY for absolute neutrophil count below 1500/mm<sup>3</sup> or neutropenic fever. Monitor blood cell counts periodically during treatment. Consider G-CSF for secondary prophylaxis. Initiate anti-infective treatment in patients with febrile neutropenia without delay.**
- **Severe diarrhea may occur. Monitor patients with diarrhea and give fluid and electrolytes as needed. At the onset of diarrhea, evaluate for infectious causes and, if negative, promptly initiate loperamide. If severe diarrhea occurs, withhold TRODELVY until resolved to ≤Grade 1 and reduce subsequent doses.**

## CONTRAINDICATIONS

- Severe hypersensitivity reaction to TRODELVY.

Please see additional Important Safety Information on next page and accompanying full [Prescribing Information](#), including **BOXED WARNING**.

**WARNINGS AND PRECAUTIONS**

**Neutropenia:** Severe, life-threatening, or fatal neutropenia can occur and may require dose modification. Neutropenia occurred in 64% of patients treated with TRODELVY. Grade 3-4 neutropenia occurred in 49% of patients. Febrile neutropenia occurred in 6%. Neutropenic colitis occurred in 1.4%. Withhold TRODELVY for absolute neutrophil count below 1500/mm<sup>3</sup> on Day 1 of any cycle or neutrophil count below 1000/mm<sup>3</sup> on Day 8 of any cycle. Withhold TRODELVY for neutropenic fever. Administer G-CSF as clinically indicated or indicated in Table 1 of USPI.

**Diarrhea:** Diarrhea occurred in 64% of all patients treated with TRODELVY. Grade 3-4 diarrhea occurred in 11% of patients. One patient had intestinal perforation following diarrhea. Diarrhea that led to dehydration and subsequent acute kidney injury occurred in 0.7% of all patients. Withhold TRODELVY for Grade 3-4 diarrhea and resume when resolved to ≤Grade 1. At onset, evaluate for infectious causes and if negative, promptly initiate loperamide, 4 mg initially followed by 2 mg with every episode of diarrhea for a maximum of 16 mg daily. Discontinue loperamide 12 hours after diarrhea resolves. Additional supportive measures (e.g., fluid and electrolyte substitution) may also be employed as clinically indicated. Patients who exhibit an excessive cholinergic response to treatment can receive appropriate premedication (e.g., atropine) for subsequent treatments.

**Hypersensitivity and Infusion-Related Reactions:** Serious hypersensitivity reactions including life-threatening anaphylactic reactions have occurred with TRODELVY. Severe signs and symptoms included cardiac arrest, hypotension, wheezing, angioedema, swelling, pneumonitis, and skin reactions. Hypersensitivity reactions within 24 hours of dosing occurred in 35% of patients. Grade 3-4 hypersensitivity occurred in 2% of patients. The incidence of hypersensitivity reactions leading to permanent discontinuation of TRODELVY was 0.2%. The incidence of anaphylactic reactions was 0.2%. Pre-infusion medication is recommended. Have medications and emergency equipment to treat such reactions available for immediate use. Observe patients closely for hypersensitivity and infusion-related reactions during each infusion and for at least 30 minutes after completion of each infusion. Permanently discontinue TRODELVY for Grade 4 infusion-related reactions.

**Nausea and Vomiting:** Nausea occurred in 64% of all patients treated with TRODELVY and Grade 3-4 nausea occurred in 3% of these patients. Vomiting occurred in 35% of patients and Grade 3-4 vomiting occurred in 2% of these patients. Premedicate with a two or three drug combination regimen (e.g., dexamethasone with either a 5-HT<sub>3</sub> receptor antagonist or an NK<sub>1</sub> receptor antagonist as well as other drugs as indicated) for prevention of chemotherapy-induced nausea and vomiting (CINV). Withhold TRODELVY doses for Grade 3 nausea or Grade 3-4 vomiting and resume with additional supportive measures when resolved to Grade ≤1. Additional antiemetics and other supportive measures may also be employed as clinically indicated. All patients should be given take-home medications with clear instructions for prevention and treatment of nausea and vomiting.

**Increased Risk of Adverse Reactions in Patients with Reduced UGT1A1 Activity:** Patients homozygous for the uridine diphosphate-glucuronosyl transferase 1A1 (UGT1A1)\*28 allele are at increased risk for neutropenia, febrile neutropenia, and anemia and may be at increased risk for other adverse reactions with TRODELVY. The incidence of Grade 3-4 neutropenia was 58% in patients homozygous for the UGT1A1\*28, 49% in patients heterozygous for the UGT1A1\*28 allele, and 43% in patients homozygous for the wild-type allele. The incidence of Grade 3-4 anemia was 21% in patients homozygous for the UGT1A1\*28 allele, 10% in patients heterozygous for the UGT1A1\*28 allele, and 9% in patients homozygous for the wild-type allele. Closely monitor patients with known reduced UGT1A1 activity for adverse reactions. Withhold or permanently discontinue TRODELVY based on clinical assessment of the onset, duration and severity of the observed adverse reactions in patients with evidence of acute early-onset or unusually severe adverse reactions, which may indicate reduced UGT1A1 function.

**Embryo-Fetal Toxicity:** Based on its mechanism of action, TRODELVY can cause teratogenicity and/or embryo-fetal lethality when administered to a pregnant woman. TRODELVY contains a genotoxic component, SN-38, and targets rapidly dividing cells. Advise pregnant women and females of reproductive potential of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with TRODELVY and for 6 months after the last dose. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with TRODELVY and for 3 months after the last dose.

**ADVERSE REACTIONS**

In the pooled safety population, the most common (≥ 25%) adverse reactions including laboratory abnormalities were decreased leukocyte count (84%), decreased neutrophil count (75%), decreased hemoglobin (69%), diarrhea (64%), nausea (64%), decreased lymphocyte count (63%), fatigue (51%), alopecia (45%), constipation (37%), increased glucose (37%), decreased albumin (35%), vomiting (35%), decreased appetite (30%), decreased creatinine clearance (28%), increased alkaline phosphatase (28%), decreased magnesium (27%), decreased potassium (26%), and decreased sodium (26%).

**In the ASCENT study** (locally advanced or metastatic triple-negative breast cancer), the most common adverse reactions (incidence ≥25%) were fatigue, diarrhea, nausea, alopecia, constipation, vomiting, abdominal pain, and decreased appetite. The most frequent serious adverse reactions (SAR) (>1%) were neutropenia (7%), diarrhea (4%), and pneumonia (3%). SAR were reported in 27% of patients, and 5% discontinued therapy due to adverse reactions. The most common Grade 3-4 lab abnormalities (incidence ≥25%) in the ASCENT study were reduced neutrophils, leukocytes, and lymphocytes.

**In the TROPiCS-02 study** (locally advanced or metastatic HR-positive, HER2-negative breast cancer), the most common adverse reactions (incidence ≥25%) were diarrhea, fatigue, nausea, alopecia, and constipation. The most frequent serious adverse reactions (SAR) (>1%) were diarrhea (5%), febrile neutropenia (4%), neutropenia (3%), abdominal pain, colitis, neutropenic colitis, pneumonia, and vomiting (each 2%). SAR were reported in 28% of patients, and 6% discontinued therapy due to adverse reactions. The most common Grade 3-4 lab abnormalities (incidence ≥25%) in the TROPiCS-02 study were reduced neutrophils and leukocytes.

**DRUG INTERACTIONS**

**UGT1A1 Inhibitors:** Concomitant administration of TRODELVY with inhibitors of UGT1A1 may increase the incidence of adverse reactions due to potential increase in systemic exposure to SN-38. Avoid administering UGT1A1 inhibitors with TRODELVY.

**UGT1A1 Inducers:** Exposure to SN-38 may be reduced in patients concomitantly receiving UGT1A1 enzyme inducers. Avoid administering UGT1A1 inducers with TRODELVY.

**Please see accompanying full [Prescribing Information](#), including **BOXED WARNING**.**